

***Arthromoniliphora araucariae* gen. & sp. nov. from Brazilian pine**

SILVANA SANTOS DA SILVA¹, LUÍS FERNANDO PASCHOLATI GUSMÃO^{1*}
& RAFAEL F. CASTAÑEDA-RUIZ²

¹*Departamento de Ciências Biológicas, Universidade Estadual de Feira de Santana,
44031-460, Feira de Santana, BA, Brazil*

²*Instituto de Investigaciones Fundamentales en Agricultura Tropical 'Alejandro de Humboldt'
(INIFAT), Académico Titular de la Academia de Ciencias de Cuba,
Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200*

*CORRESPONDENCE TO: lgusmao@uefs.br

ABSTRACT—A new genus and species, *Arthromoniliphora araucariae*, is described and illustrated. This fungus is distinguished by differentiated hyaline conidiophores that are branched above the globose base and robustly cylindrical below, moniliform and subulate toward the apex. The conidia are thallic-arthric, cylindrical, bacilliform to cuneate, and hyaline and occur in unbranched chains formed by disarticulation of the moniliform, subulate conidiophore branches.

KEY WORDS—*Araucaria angustifolia*, asexual fungi, conidial fungi, taxonomy

Introduction

During a survey of hyphomycetes associated with plant litter in Floresta Nacional de São Francisco de Paula, Rio Grande do Sul State, Brazil, an interesting fungus was collected. This fungus shows remarkable differences from all previously described genera (Seifert et al. 2011) and is therefore described as a new genus and species.

Material & methods

Samples of litter of *Araucaria angustifolia* (Brazilian pine; Paraná pine) were placed in paper bags. In the laboratory the samples were placed in Petri dish moist chambers and stored in a 170 L polystyrene box with 200 mL sterile water plus 2 mL glycerol at 25°C for 30 days (Castañeda-Ruiz 2005). Mounts were prepared in PVL (polyvinyl

alcohol, lactic acid, and phenol). Measurements were made at a magnification of $\times 1000$. Micrographs were obtained with an Olympus BX51 microscope equipped with bright field and Nomarski interference optics. The type specimen is deposited in the Herbarium of Universidade Estadual de Feira de Santana, Bahia, Brazil (HUEFS).

Taxonomy

Arthromoniliphora S.S. Silva, Gusmão & R.F. Castañeda, **gen. nov.**

MYCOBANK MB 813427

Differs from *Arthrographis* and *Geotrichum* by its moniliform, closely fasciculate conidiophores and its conidia with a highly variable number of septa.

TYPE SPECIES: *Arthromoniliphora araucariae* S.S. Silva et al.

ETYMOLOGY: Greek *Arthro-*, meaning jointed (referring to thallic disarticulation) + *-monili-* Latin, meaning cylindrical but contracted at regular intervals + *-phora* referring to the conidiophores.

Conidial fungi. COLONIES on the natural substratum effuse, white. CONIDIOPHORES differentiated, single, densely fasciculate, branched, moniliform, septate, hyaline. CONIDIOGENOUS HYPHAE moniliform, subulate, holothallic, discrete, determinate. CONIDIAL SECESSION schizolytic. CONIDIA thallic-arthric, in unbranched chains, unpigmented, unicellular or multiseptate, cylindrical, doliiform or cuneate, forming by disarticulation of the terminal moniliform conidiogenous conidiophore branches.

Arthromoniliphora araucariae S.S. Silva, Gusmão & R.F. Castañeda, **sp. nov.**

MYCOBANK MB 813428

FIGS 1–3

Differs from all *Arthrographis* and *Geotrichum* spp. by its densely fasciculate conidiophores that are moniliform and cylindrical below and subulate above, and its cylindrical, unicellular to multiseptate conidia.

TYPE: Brazil, Rio Grande do Sul State, São Francisco de Paula, Floresta Nacional de São Francisco de Paula, 29°25'S 50°23'W, alt. 838 m, on decaying twig of *Araucaria angustifolia* (Bertol.) Kuntze (*Araucariaceae*), 10.III 2015, S.S. Silva (Holotype: HUEFS 216008).

ETYMOLOGY: Latin *araucariae*, meaning growth on an *Araucaria* species.

COLONIES on the natural substratum effuse, hairy, white. Mycelium superficial and immersed. Hyphae septate, branched, smooth, hyaline, 3–5 μm diam, sometimes forming a rudimentary globulose stroma. CONIDIOPHORES distinct, single, erect, compact fasciculate, branched just at one level near the base, moniliform, subulate toward apex of each branch, hyaline, multiseptate, 60–250 $\times$ 5–6 μm . CONIDIOGENOUS BRANCHES moniliform, subulate, holothallic, multiseptate, hyaline, discrete, determinate, 38–100 $\times$ 2.5–4 μm . CONIDIAL SECESSION schizolytic. CONIDIA thallic-arthric, in unbranched chains, hyaline, 0–36-septate, cylindrical, doliiform or cuneate, truncate at




FIG. 1. *Arthromoniliphora araucariae* (ex holotype, HUEFS 216008). A, B. Conidiophores, conidiogenous branches, and conidia. C. Rudimentary stroma and basal part of conidiophores.

the ends, sometimes truncate at the base and obtuse at the apex, $3.5\text{--}92 \times 2.5\text{--}4 \mu\text{m}$ forming by disarticulation of the terminal, subuliform, moniliform conidiogenous branches of the conidiophores




FIG. 2. *Arthromoniliphora araucariae* (ex holotype, HUEFS 216008).
Conidial chains and conidia.

COMMENTS—The genera *Arthrographis* G. Cochet ex Sigler & J.W. Carmich., *Geomyces* Traaen, and *Geotrichum* Link superficially resemble *Arthromoniliphora* in their hyaline thallic-arthric conidia produced by disarticulation of fertile hyphae. In *Arthrographis*, however, conidiophores are multi-branched, producing




FIG. 3. *Arthromoniliphora araucariae* (ex holotype, HUEFS 216008).
A. Conidiophores, conidiogenous branches, and conidia. B. Conidia.

conidia that are globose, cylindrical, Y-shaped or irregular (usually discoid), rounded or subtruncate at the ends after disarticulation (Sigler & Carmichael 1976, Giraldo et al 2014). *Geomyces* has branched hyaline conidiophores, but the conidia have separating cells and the conidial secession is rhexolytic. *Geotrichum* is characterized by undifferentiated conidiophores, and after disarticulation of the fertile hyphae the conidia are unicellular, cylindrical, Y-shaped to irregular, sometimes forming chains, or slimy or yeast-like (Cole & Kendrick 1969, Sigler & Carmichael 1976, Hoog et al. 2011, Seifert et al. 2011). *Theodgonia* B. Sutton (Sutton, 1973) is somewhat similar to *Arthromoniliphora*, but its conidiogenous cells were described as polyblastic and sympodial and its conidia as blastocatenate, cylindrical, and 0–3-septate by Sutton (1973), while Seifert et al. (2011) described the conidial ontogeny as thallic-arthric. The *Theodgonia* conidiophores are branched and cylindrical but not moniliform and subulate as in *Arthromoniliphora*

Acknowledgments

The authors express their sincere gratitude to Dr. De-Wei Li and Prof Xiu-Guo Zhang for their critical review of the manuscript. The authors thank the National Council for Scientific and Technological Development (CNPq) (Proc. 142014/2011–7) and “Programa de Pesquisa em Biodiversidade do Semiárido (PPBio Semiárido -CNPq/MCTI) for supporting this study. RFCR is grateful to Cuban Ministry of Agriculture and ‘Programa de Salud Animal y Vegetal’, project P131LH003033 Cuban for facilities. We acknowledge the assistance provided by Dr. P.M. Kirk and Drs. V. Robert and G. Stegehuis through the Index Fungorum and MycoBank websites. Dr. Lorelei Norvell’s editorial and Dr. Shaun Pennycook’s nomenclatural reviews are greatly appreciated.

Literature cited

- Castañeda Ruiz RF. 2005. Metodología en el estudio de los hongos anamorfos. 182–183 in: Anais do V Congresso Latino Americano de Micologia. Brasilia.
- Cole GT, Kendrick WB. 1969. Conidium ontogeny in hyphomycetes. The arthrospores of *Oidiodendron* and *Geotrichum*, and the endoarthrospores of *Sporendonema*. Canadian Journal of Botany 47: 1773–1780. <https://doi.org/10.1139/b69-255>
- Giraldo A, Gené J, Sutton DA, Madrid H, Cano J, Crous PW, Guarro J. .2014. Phylogenetic circumscription of *Arthrographis* (Eremomycetaceae, Dothideomycetes) Persoonia 32: 102–114. <https://doi.org/10.3767/003158514X680207>
- Hoog GS de, Guarro J, Gené J, Figueras MJ.2011. Atlas of clinical fungi, electronic version 3.1. CBS-KNAW Fungal Biodiversity Centre, Utrecht, Netherlands
- Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. CBS Biodiversity Series 9. 997 p.
- Sigler L, Carmichael JW. 1976. Taxonomy of *Malbranchea* and some other hyphomycetes with arthroconidia. Mycotaxon 4: 349–388.
- Sutton BC. 1973. Some hyphomycetes with holoblastic sympodial conidiogenous cells. Transactions of the British Mycological Society 61: 417–429. [https://doi.org/10.1016/S0007-1536\(73\)80112-3](https://doi.org/10.1016/S0007-1536(73)80112-3)